

Document details

IOP Conference Series: Materials Science and Engineering
Volume 342, Issue 1, 6 April 2018, Article number 012017
International Conference on Innovative Technology, Engineering and Sciences 2018, iCITES 2018; Universiti Malaysia Pahang (UMP) Pekan Campus LibraryPekan, Pahang; Malaysia; 1 March 2018 through 2 March 2018; Code 135946

The classification of hunger behaviour of Lates Calcarifer through the integration of image processing technique and k-Nearest Neighbour learning algorithm (Conference Paper) [\(Open Access\)](#)

Taha, Z.^a ✉, Razman, M.A.M.^a, Abdul Ghani, A.S.^a, Abdul Majeed, A.P.P.^a, Musa, R.M.^a, Adnan, F.A.^a, Sallehudin, M.F.^b, Mukai, Y.^b

^aInnovative Manufacturing Mechatronics and Sports Lab (IMAMS), Faculty of Manufacturing Engineering, Universiti Malaysia Pahang, Pekan Campus, Pekan, Pahang, Malaysia

^bKulliyyah of Science, International Islamic University Malaysia, Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, Pahang, Kuantan, Malaysia

Abstract

⌵ View references (14)

Fish Hunger behaviour is essential in determining the fish feeding routine, particularly for fish farmers. The inability to provide accurate feeding routines (under-feeding or over-feeding) may lead the death of the fish and consequently inhibits the quantity of the fish produced. Moreover, the excessive food that is not consumed by the fish will be dissolved in the water and accordingly reduce the water quality through the reduction of oxygen quantity. This problem also leads the death of the fish or even spur fish diseases. In the present study, a correlation of Barramundi fish-school behaviour with hunger condition through the hybrid data integration of image processing technique is established. The behaviour is clustered with respect to the position of the school size as well as the school density of the fish before feeding, during feeding and after feeding. The clustered fish behaviour is then classified through k-Nearest Neighbour (k-NN) learning algorithm . Three different variations of the algorithm namely cosine, cubic and weighted are assessed on its ability to classify the aforementioned fish hunger behaviour . It was found from the study that the weighted k-NN variation provides the best classification with an accuracy of 86.5%. Therefore, it could be concluded that the proposed integration technique may assist fish farmers in ascertaining fish feeding routine. © Published under licence by IOP Publishing Ltd.

Indexed keywords

Engineering controlled terms:

- Agriculture
- Data integration
- Dissolved oxygen
- Feeding
- Fish
- Image classification
- Nearest neighbor search
- Water quality

Engineering uncontrolled terms

- Fish behaviours
- Fish disease
- Hybrid datum
- Integration of images
- Integration techniques
- K nearest neighbours (k-NN)
- K-nearest neighbours
- Reduction of oxygen

Engineering main heading:

- Learning algorithms

Funding details

Funding number	Funding sponsor	Acronym	Funding opportunities
	Universiti Malaysia Pahang	UMP	

Metrics ⓘ

- 0 Citations in Scopus
- 0 Field-Weighted Citation Impact



PlumX Metrics ⌵
Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

Related documents

- The Identification of Hunger Behaviour of Lates Calcarifer through the Integration of Image Processing Technique and Support Vector Machine
Taha, Z. , Razman, M.A.M. , Adnan, F.A.
(2018) *IOP Conference Series: Materials Science and Engineering*
- The Identification of hunger behaviour of lates calcarifer using k-nearest neighbour
Taha, Z. , Razman, M.A.M. , Adnan, F.A.
(2018) *Lecture Notes in Mechanical Engineering*
- Cave-dwelling fish provide clues to the circadian cycle
Mejia, R.
(2011) *PLoS Biology*

View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

ISSN: 17578981

Source Type: Conference
Proceeding

Original language: English

DOI: 10.1088/1757-899X/342/1/012017



Document Type: Conference Paper

Sponsors: Intel Malaysia,SCANIA Malaysia,Tourism
Pahang

Publisher: Institute of Physics Publishing

References (14)

[View in search results format >](#)

☐ All [Export](#)  [Print](#)  [E-mail](#) [Save to PDF](#) [Create bibliography](#)

- ☐ 1 Cavallari, N., Frigato, E., Vallone, D., Fröhlich, N., Lopez-Olmeda, J.F., Foà, A., Berti, R., (...), Foulkes, N.S.
A blind circadian clock in cavefish reveals that opsins mediate peripheral clock
photoreception ([Open Access](#))

(2011) *PLoS Biology*, 9 (9), art. no. e1001142. Cited 99 times.
<http://www.plosbiology.org/article/fetchObjectAttachment.action?uri=info%3Adoi%2F10.1371%2Fjournal.pbio.1001142&representation=PDF>
doi: 10.1371/journal.pbio.1001142

[View at Publisher](#)

- ☐ 2 Hansen, M.J., Schaerf, T.M., Ward, A.J.W.
The effect of hunger on the exploratory behaviour of shoals of mosquitofish *Gambusia holbrooki*

(2015) *Behaviour*, 152 (12-13), pp. 1659-1677. Cited 14 times.
<http://www.brill.com/behaviour>
doi: 10.1163/1568539X-00003298

[View at Publisher](#)

- ☐ 3 Beale, A., Guibal, C., Tamai, T.K., Klotz, L., Cowen, S., Peyric, E., Reynoso, V.H., (...), Whitmore, D.
(2013) *Circadian Rhythms in Mexican Blind Cavefish *Astyanax Mexicanus* in the Lab and in the Field*, 4, p. 2769. Cited 2 times.

- ☐ 4 Priyadarshana, T., Asaeda, T., Manatunge, J.
Hunger-induced foraging behavior of two cyprinid fish: *Pseudorasbora parva* and *Rasbora daniconius*

(2006) *Hydrobiologia*, 568 (1), pp. 341-352. Cited 13 times.
doi: 10.1007/s10750-006-0201-5

[View at Publisher](#)

- ☐ 5 Chapman, B.B., Morrell, L.J., Krause, J.
Unpredictability in food supply during early life influences boldness in fish

(2010) *Behavioral Ecology*, 21 (3), pp. 501-506. Cited 49 times.
doi: 10.1093/beheco/arq003

[View at Publisher](#)

-
- ☐ 6 Sanchez-Vázquez, F.J., Madrid, J.A., Zamora, S.
Circadian Rhythms of Feeding Activity in Sea Bass, *Dicentrarchus labrax* L.: Dual Phasing Capacity of Diel Demand-Feeding Pattern

(1995) *Journal of Biological Rhythms*, 10 (3), pp. 256-266. Cited 112 times.
doi: 10.1177/074873049501000308

[View at Publisher](#)
-
- ☐ 7 Biswas, G., Thirunavukkarasu, A.R., Sundaray, J.K., Kailasam, M.
Optimization of feeding frequency of Asian seabass (*Lates calcarifer*) fry reared in net cages under brackishwater environment

(2010) *Aquaculture*, 305 (1-4), pp. 26-31. Cited 31 times.
doi: 10.1016/j.aquaculture.2010.04.002

[View at Publisher](#)
-
- ☐ 8 Harpaz, S., Hakim, Y., Barki, A., Karplus, I., Slosman, T., Eroldogan, O.T.
Effects of different feeding levels during day and/or night on growth and brush-border enzyme activity in juvenile *Lates calcarifer* reared in freshwater re-circulating tanks

(2005) *Aquaculture*, 248 (1-4), pp. 325-335. Cited 30 times.
doi: 10.1016/j.aquaculture.2005.04.033

[View at Publisher](#)
-
- ☐ 9 Ariyomo, T.O., Watt, P.J.
Effect of hunger level and time of day on boldness and aggression in the zebrafish *Danio rerio*

(2015) *Journal of Fish Biology*, 86 (6), pp. 1852-1859. Cited 12 times.
doi: 10.1111/jfb.12674

[View at Publisher](#)
-
- ☐ 10 Josep, A.
Repeatability of circadian behavioural variation revealed in free-ranging marine fish
(2017) *Royal Society Publishing*, 4, p. 11. Cited 2 times.
M M-B a A C-C
-
- ☐ 11 Cortes, C., Vapnik, V.
Support-Vector Networks

(1995) *Machine Learning*, 20 (3), pp. 273-297. Cited 17740 times.
doi: 10.1023/A:1022627411411

[View at Publisher](#)
-
- ☐ 12 Dutta, M.K., Sengar, N., Kamble, N., Banerjee, K., Minhas, N., Sarkar, B.
Image processing based technique for classification of fish quality after cypermethrine exposure

(2016) *LWT - Food Science and Technology*, 68, pp. 408-417. Cited 8 times.
<http://www.elsevier.com/inca/publications/store/6/2/2/9/1/0/index.htm>
doi: 10.1016/j.lwt.2015.11.059

[View at Publisher](#)
-

13

Cubitt, K.F., Williams, H.T., Rowsell, D., McFarlane, W.J., Gosine, R.G., Butterworth, K.G., McKinley, R.S.
Development of an intelligent reasoning system to distinguish hunger states in Rainbow trout (*Oncorhynchus mykiss*)

(2008) *Computers and Electronics in Agriculture*, 62 (1), pp. 29-34. Cited 8 times.
doi: 10.1016/j.compag.2007.08.010

View at Publisher

14

Taha, Z., Musa, R.M., P.P. Abdul Majeed, A., Alim, M.M., Abdullah, M.R.
The identification of high potential archers based on fitness and motor ability variables: A Support Vector Machine approach

(2018) *Human Movement Science*, 57, pp. 184-193. Cited 6 times.
<http://www.elsevier.com/locate/humov>
doi: 10.1016/j.humov.2017.12.008

View at Publisher

© Copyright 2018 Elsevier B.V., All rights reserved.

- About Scopus
- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

- Language
- 日本語に切り替える
- 切换到简体中文
- 切換到繁體中文
- Русский язык

- Customer Service
- Help
- Contact us